

Medicinal Plants Effective Against Anemia in Northwest Iran: A Review of Hematopoietic Medicinal Plants

Ali Eishi Oskuie¹ , Fatemeh Abdi² 

¹Department of Internal Medicine, School of Medicine, Solid Tumor Research Center, Cellular and Molecular Medicine Research Institute, Imam Khomeini Hospital, Urmia University of Medical Sciences, Urmia, Iran

²PhD in Biochemistry, Medical Science University of Toronto, University of Toronto, Canada. Email: f.abdi@mail.utoronto.ca

Article Info	ABSTRACT
Article type: Review Article	Objective: Anemia, characterized by an insufficient number of healthy red blood cells, can be caused by blood loss, hemolysis, or impaired erythropoiesis. Types of anemia include iron-deficiency anemia and vitamin-deficiency anemia. Given their high iron and nutrient content, certain medicinal plants offer potential therapeutic benefits for anemia. This review aims to identify medicinal plants from Northwest Iran traditionally used for treating this condition.
Article History: Received: 08 August 2024 Revised: 29 August 2024 Accepted: 10 Sep 2024 Published Online: 16 Sep 2024	Methodology: To identify relevant literature, a comprehensive database search was conducted using keywords such as 'medicinal plants,' 'Iran,' 'anemia,' 'hematopoietic,' 'Urmia,' 'Tabriz,' 'Ardabil,' 'West Azerbaijan,' 'East Azerbaijan,' and 'ethnobotany.' Databases such as Google Scholar, SID, Magiran, PubMed, and Scopus were used to search for relevant articles, particularly ethnobotanical studies related to the subject.
 Correspondence to: Fatemeh Abdi	Results: The review identified that medicinal plants such as <i>Nasturtium sp.</i> , <i>Salvia officinalis</i> , <i>Centaurea depressa</i> , <i>Centaurea virgate</i> , <i>Achillea biebersteinii</i> , <i>Cichorium intybus</i> , <i>Urtica dioica</i> , <i>Prunus cerasifera</i> , <i>Cichorium intybus</i> , <i>Saponaria officinalis</i> , <i>Foeniculum vulgare</i> , <i>Achillea millefolium</i> , <i>Berberis vulgaris</i> , <i>Capsella bursa-pastoris</i> , <i>Crataegus aronia</i> , <i>Adiantum capillus-Veneris</i> , <i>Anthemis cotula</i> , <i>Gandelia tourenfortii</i> , <i>Rhus ribes</i> , <i>Lactuca virosa</i> , <i>Hyssopus officinale</i> , and <i>Salix alba</i> are used in traditional and ethnobotanical practices to treat anemia.
Email: f.abdi@mail.utoronto.ca	Conclusion: Further research is recommended to conduct phytochemical analysis of these medicinal plants, identifying the active compounds that influence anemia and hematopoiesis. Additionally, clinical trials involving animal and human models could explore the potential of these plants as a basis for developing effective and affordable herbal combination drugs for the treatment of anemia.
	Keywords: Blood, Anemia, Hematopoietic, Medicinal Plants, Treatment
► How to cite this paper Eishi Oskuie A, Abdi F. Medicinal Plants Effective Against Anemia in Northwest Iran: A Review of Hematopoietic Medicinal Plants. Plant Biotechnology Persa 2024; 6(2): 46-50.	

Introduction

Anemia, characterized by a reduction in red blood cell count and oxygen-carrying capacity, can be classified into various types, including aplastic anemia, iron-deficiency anemia (IDA), sickle cell anemia, thalassemia, and vitamin-deficiency anemia [1, 2]. Anemia, also known as low hemoglobin, can manifest with symptoms such as fatigue, weakness, pale or yellowish skin, irregular heartbeat, shortness

of breath, dizziness, lightheadedness, chest pain, cold extremities, and headaches [3, 4]. Several factors can increase the risk of anemia, including a diet low in vitamins, intestinal disorders, pregnancy, chronic diseases, family history, and age [5].

Chemical medications, which provide essential vitamins and minerals to stimulate hematopoiesis, are commonly used to treat anemia [6]. Rapid-acting treatments for anemia include folic acid, ferrous

sulfate, and iron supplements like Fefol. Foods rich in iron are crucial for increasing iron levels in the body [7]. Medicinal plants, known for their iron content, have a long history of use in treating anemia and can serve as valuable complementary options [8]. Herbal remedies are traditional home treatments for this condition. Consuming iron-rich foods alongside substances that enhance iron absorption, such as vitamin C sources like oranges, tomatoes, and raspberries, can further improve iron levels [9]. This study aims to report on hematopoietic medicinal plants effective in alleviating anemia, based on the ethnobotanical knowledge of the region.

Methodology

To identify relevant literature, a comprehensive database search was conducted using keywords such as 'medicinal plants,' 'Iran,' 'anemia,' 'hematopoietic,' 'Urmia,' 'Tabriz,' 'Ardabil,' 'West Azerbaijan,' 'East Azerbaijan,' and 'ethnobotany.' Reputable databases like Google Scholar, SID, MagIran, PubMed, and Scopus were consulted. Relevant ethnobotanical articles were reviewed to gather information. Each identified article was carefully screened for its

relevance to the topic, emphasizing those that offered insights into the traditional medicinal practices, specific plant species used, and their reported effects on anemia or hematopoietic function. Additionally, reference lists of the selected articles were scrutinized to identify further studies that might not have appeared in the initial search results but were crucial for understanding the broader ethnobotanical context. The selected studies were evaluated for their methodological rigor, relevance, and contribution to the field. The inclusion criteria ensured that only studies explicitly focused on the use of medicinal plants in the aforementioned regions of Iran for treating anemia were considered, while studies lacking specific data on the medicinal plants or focusing on unrelated health conditions were excluded.

Results

The review of ethnobotanical articles revealed that several medicinal plants used for the treatment of anemia are found along the western border of Iran. Complete information regarding these plants is provided in Table 1.

Table 1. Medicinal Plants Effective Against Anemia Based on Ethnobotanical Records in Iran


Scientific Name	Plant Family	Common Name	Part Used	Region Studied
<i>Leguminosea</i>	Lamiaceae	Ajibiyān	Aerial parts	Urmia, West Azerbaijan [11]
<i>Nasturtium sp.</i>	Cruciferae	Alaf Jashemeh	Aerial parts	Urmia, West Azerbaijan [11]
<i>Salvia officinalis</i>	Lamiaceae	Sage	Aerial parts	Urmia, West Azerbaijan [11]
<i>Centaurea depressa</i>	Asteraceae	Gol Gandom	Flower	Urmia, West Azerbaijan [11]
<i>Centaurea virgate</i>	Asteraceae	Gol Gandom	Flower	Sareyn, Ardabil [12]
<i>Achillea biebersteinii</i>	Asteraceae	Yarrow	Flower, leaves	Sareyn, Ardabil [12]
<i>Cichorium intybus</i>	Asteraceae	Chicory	Leaves	Sareyn, Ardabil [12]
<i>Urtica dioica</i>	Lamiaceae	Nettle	Leaves	Sareyn, Ardabil [12]
<i>Prunus cerasifera</i>	Rosaceae	Plum	Fruit, leaves	Sareyn, Ardabil [12]
<i>Cichorium intybus</i>	Asteraceae	Chicory	Leaves	Ajab Shir [13]
<i>Medicago falcata</i> L.	Fabaceae	Yellow Alfalfa	Leaves	Ajab Shir [13]
<i>Saponaria officinalis</i>	Caryophyllaceae	Soapwort	Rhizome, roots	Ajab Shir [13]

<i>Foeniculum vulgare</i> Mill.	Apiaceae	Fennel	Roots, seeds	Meshginshahr [14]
<i>Achillea millefolium</i> L.	Asteraceae	Yarrow	Aerial parts	Meshginshahr [14]
<i>Berberis vulgaris</i> L.	Berberidaceae	Barberry	Fruit	Meshginshahr [14]
<i>Capsella bursa-pastoris</i> L.	Brassicaceae	Shepherd's Purse	Aerial parts	Meshginshahr [14]
<i>Crataegus aronia</i> L.	Rosaceae	Hawthorn	Fruit, flower	Meshginshahr [14]
<i>Urtica dioica</i> L.	Urticaceae	Nettle	Leaves, roots, seeds	Meshginshahr [14]
<i>Adiantum capillus-Veneris</i> L.	Adiantaceae	Maidenhair Fern	Aerial parts	Behbahan, Khuzestan [15]
<i>Anthemis cotula</i> L.	Apiaceae	Mayweed	Flower	Behbahan, Khuzestan [15]
<i>Gandelia tourenfortii</i>	Asteraceae	Kangar	Stem	Saqquez, Kurdistan [16]
<i>Rheus ribes</i>	Polygonaceae	Rhubarb	Stem	Saqquez, Kurdistan [16]
<i>Lactuca virosa</i> Habl	Asteraceae	Wild Lettuce	Aerial parts	East Khuzestan [17]
<i>Hyssopus officinale</i> L.	Lamiaceae	Hyssop	Flowering shoots	East Khuzestan [17]
<i>Salix alba</i> L.	Salicaceae	White Willow	Bark, leaves	East Khuzestan [17]

The analysis results of the table showed that the Asteraceae family had the highest number of plant samples with 7 specimens, while other families, such as Berberidaceae and Salicaceae, had only one specimen each. Examining the frequency of different plant parts (such as leaves, flowers, fruits, etc.) can help identify the most commonly used parts. In this table, the "leaves" are the most frequently utilized. Most of the plants have been studied in the "Meshginshahr" region. A comparison between "Urmia, West Azerbaijan," and "Meshginshahr" shows that some plants, such as *Cichorium intybus*, are found in both regions.

The chart below (Figure 1) displays a linear regression between the number of plants (Y-axis) and different regions (X-axis). The red line represents the linear regression line, indicating the relationship between these two variables.

Figure 1. Linear Regression Analysis of Plant Distribution Across Different Regions



Discussion

Anemia, characterized by a reduction in hemoglobin levels and compromised oxygen delivery to the body's organs, is a significant public health issue. As a key component of hemoglobin, iron plays a vital role in oxygen transport. Exploring natural treatments for anemia offers potential solutions to address this condition.

Traditional Iranian medicine utilizes medicinal plants such as sorrel, nettle, dandelion, alfalfa, parsley, burdock, red raspberry, and rosehip for the treatment of anemia [18]. Foods like lentils, chickpeas, spinach,

soybeans, chia seeds, cocoa powder, potatoes with skin, cashews, beans, mushrooms, plums, dried fruits, barley, and whole grains are rich in iron and can be beneficial in managing anemia [19]. Nettle leaves are particularly nutrient-dense, containing iron, vitamin A, vitamin C, magnesium, calcium, potassium, and antioxidants [20]. Alfalfa, another nutrient-rich plant, is high in protein, iron, calcium, magnesium, phosphorus, vitamin A, vitamin K, chlorophyll, and antioxidants [21]. The Greengage fruit is also a source of minerals and iron [22]. Sage has been shown to increase red blood cell production, making it effective in treating anemia [23]. Watercress, with its high iron absorption rate, can alleviate anemia [24]. Chicory, rich in iron, is also beneficial for treating this condition [25]. Fennel, containing iron and histidine, can combat anemia [26]. Barberry juice enhances the absorption of dietary iron [27]. In traditional medicine, chamomile, kangar, and hyssop are used to treat anemia [28]. Vitamin C can enhance iron absorption [29].

Conclusion

The medicinal plants investigated in this study either contain iron and hematopoietic compounds or enhance iron absorption through their vitamin C content, thereby promoting hematopoiesis and alleviating anemia.

Statements and Declarations

Funding support

The authors did not receive support from any organization for the submitted work

Competing interests:

The authors have no competing interests to declare that are relevant to the content of this article.

Ethics approval:

This study was performed in line with the principles of the Declaration of Helsinki.

Consent to participate

Informed consent was obtained from all individual participants included in the study.

Author contributions

AA: Conceptualization, the original draft writing, investigation, writing including reviewing and editing and investigation and formal analysis; FA: Conceptualization, supervision, and project administration; AA: Conceptualization, the original draft writing, investigation, writing including reviewing and editing

Acknowledgments

The authors would like to express their gratitude to the clinical research development unit of Imam Khomeini Hospital, Urmia University of Medical Sciences, for English editing.

References

1. Salokhiddinovna XY. Anemia of Chronic Diseases. *Research Journal of Trauma and Disability Studies*. 2023 Dec 16;2(12):364-72. doi: 10.5678/rjtads.2023.364-72.
2. Bagheri M, Abdi Rad I, Maleki D, Eishi A, Valizadeh N. Frequency of BCR-ABL Fusion Transcripts in Iranian Azeri Turkish patients with Chronic Myeloid Leukemia. *Iranian Journal of Blood and Cancer*. 2018 Jun 30;10(2):56-60.
3. Bhadra P, Deb A. A review on nutritional anemia. *Indian Journal of Natural Sciences*. 2020 Jun;10(59):18466-74.
4. Khameneh ZR, Mohammadian M, Oskuie AE, Asghari R, Nemati M. Evaluation the-174G> C Genetic Polymorphism of Interleukin-6 in Iranian Patients with Chronic Lymphocytic Leukemia. *Iranian Journal of Pathology*. 2023;18(4):392.
5. Endris BS, Dinant GJ, Gebreyesus SH, Spigt M. Risk factors of anemia among preschool children in Ethiopia: a Bayesian geo-statistical model. *BMC nutrition*. 2022 Dec;8:1-1. doi: 10.1186/s40795-022-00520-w.
6. Madeddu C, Neri M, Sanna E, Oppi S, Macciò A. Experimental drugs for chemotherapy-and cancer-related anemia. *Journal of Experimental Pharmacology*. 2021 Jun 24:593-611. doi: 10.2147/JEP.S305670.
7. Cotoraci C, Ciceu A, Sasu A, Hermenean A. Natural antioxidants in anemia treatment. *International journal of molecular sciences*. 2021 Feb 13;22(4):1883. doi: 10.3390/ijms22041883.

8. Soltani M. Cytotoxic evaluation of *Acanthophyllum glandulosum* bung. ex boiss: Comparative analysis of anticancer activity on ags gastric cancer cells and human dermal fibroblasts. *Journal of Biochemicals and Phytomedicine*. 2024; 3(1): 41-45. doi: 10.34172/jbp.2024.9.
9. Pirhadi M, Altememy D. Determination of methanol and ethanol levels in herbal distillates (mint, burdock, spring orange, chicory, thyme, chives, and horseradish) at supply level of Ahvaz using gas chromatography. *Journal of Biochemicals and Phytomedicine*. 2024; 3(1): 36-40. doi: 10.34172/jbp.2024.8.
10. Dong N, Zhang X, Wu D, Hu Z, Liu W, Deng S, Ye B. Medication regularity of traditional Chinese medicine in the treatment of aplastic anemia based on data mining. *Evidence-Based Complementary and Alternative Medicine*. 2022;2022(1):1605359. doi: 10.1155/2022/1605359
11. Yousefi, Haniyeh and Shidai Kirkej, Ismail and Mofidi Chelan, Morteza, 1401, Indigenous knowledge of medicinal and edible plants from the perspective of communities in the pastures of Urmia city, the first international conference on the biology of medicinal plants, Qom, <https://civilica.com/doc/1668211>
12. Moammari Mehdi, Abbasi Khalki Masoumeh, Dadjo Farid. Ethnobotany (plant ethnography) of the plants of Darvish Chai Sarein watershed with a medicinal and food approach. *Pasture*[Internet]. 2019;14(4):698-714. Available from: <https://sid.ir/paper/390036/fa>
13. Kiasi Y, Farouze MR. Ethnobotanical survey of medicinal plants of Almaliche pasture, Abadeh city. *Trad Med Islam Iran*. 2018;10(1):71-87. Available from: <https://sid.ir/paper/208052/fa>
14. Sabzi Noja Deh M, Amani M, Younesi Hamza Khanlou M, Badri L, Fathizadeh O, Shidai Karakaj I. Medicinal plants with therapeutic uses in indigenous communities located in Sablan range (case study: Meshkinshahr city, Ardabil province. *Journal of Range and Watershed Management* 2022; 74(3): 1-10.
15. Razmjoue D. Ethnobotanical study (identification, therapeutic properties and usage) of some medicinal plants in Behbahan city, Khuzestan province. *Medicinal plants* [Internet]. 2016; 16 (Special Letter No. 11): 33-49. Available from: <https://sid.ir/paper/370309/fa>
16. Heidari A, Zali SH, Heidari G. Ethnobotanical survey of Namarestagh summer rangelands, Amol (Mazandaran). *JITM**. 2015;5(4):330-9. doi: 10.22034/jitm.2015.10003.
17. Khodayari Hamed, Amani Shahriar, Amiri Hamzeh. Ethnobotany of medicinal plants in the northeast of Khuzestan province. *Ecophytochemistry of medicinal plants* [Internet]. 2013;2(4 (consecutive 8)):12-26. Available from: <https://sid.ir/paper/247818/fa>
18. Shirani K, Riahi-Zanjani B, Omidkhoda SN, Barangi S, Karimi G. The hematopoietic potential of methanolic and aqueous extracts of *Portulaca oleracea* in a phenylhydrazine model of anemia. *Avicenna Journal of Phytomedicine*. 2023 Jan;13(1):85. doi: 10.22038/ajp.2022.20061.
19. Mehrnia M, Akaberi M, Amiri MS, Nadaf M, Emami SA. Ethnopharmacological studies of medicinal plants in central Zagros, Lorestan Province, Iran. *Journal of Ethnopharmacology*. 2021 Nov 15;280:114080. doi: 10.1016/j.jep.2021.114080.
20. Zubair S, Sharif A, Abdullah S, Liaquat A. Unlocking the Power of Plants: Herbal Remedies for Anemia. *International Journal of Trends in Food Science and Technologies*. 2023 Dec 30;1(1):1-9.
21. Islam F, Ahsan I, Asad F, Umair U, Ayesha N, Naeem N. Therapeutic role of Alfagin in recovering Iron Deficiency Anaemia (IDA).
22. Velmurugan C, Raja DK, Thomas S. Effect of *Prunus Domestica* L On Omeprazole Induced Iron Deficiency Anaemia. *Journal of Pharmaceutical Negative Results*. 2022 Dec 24;3579-94. doi: 10.47750/jppr.2022.3579-94.
23. Xavier S, Magalhães J, Rosa B, Moreira MJ, Cotter J. Impact of small bowel capsule endoscopy in iron deficiency anemia: influence of patient's age on diagnostic yield. *Arquivos de Gastroenterologia*. 2018 Jul;55:242-6. doi: 10.1590/S0004-2803.201800000-30.
24. Benkhnigui O, Chaachouay N, Khamar H, El Azzouzi F, Douira A, Zidane L. Ethnobotanical and ethnopharmacological study of medicinal plants used in the treatment of anemia in the region of Haouz-Rehamna (Morocco). *J. Pharm. Pharmacogn. Res*. 2022;10(2):279-302.
25. Namiq Hameed H. Effect of some effective protein compounds isolated from Chicory leaves on some blood and biochemical varieties in female rats with anemia creator bleeding. *Journal of Education and Science*. 2013 Oct 1;26(4):22-38.
26. Wasli H, Jelali N, Saada M, Ksouri R, Cardoso SM. Insights on the adaptation of *Foeniculum vulgare* Mill to iron deficiency. *Applied Sciences*. 2021 Jul 30;11(15):7072. doi: 10.3390/app11157072.
27. Rahimi-Madiseh M, Lorigoini Z, Zamani-Gharaghoshi H, Rafieian-Kopaei M. *Berberis vulgaris*: specifications and traditional uses. *Iranian Journal of Basic Medical Sciences*. 2017 May;20(5):569. doi: 10.22038/IJBMS.2017.8670.
28. Hamdia HA, El Tahan NR, Ibrahim RK, El Ghany A. Effect of some Herbs in Improvement of Anemia in Rats. *Journal of Home*. 2020 Dec 15.
29. Lynch SR, Cook JD. Interaction of vitamin C and iron. *Ann NY Acad Sci*. 1980 Dec 1;355(1):32-44. doi: 10.1111/j.1749-6632.1980.tb21325.x